

## Worksheet 1.3: Gröbner Bases Pt. I

Let  $R$  be any ring. In this course, “ring” means *commutative ring with unity*. Throughout this worksheet we fix a field  $\mathbb{K}$ .

---

**Polynomial Division.** Recall the following classical theorem:

**Theorem 1.** *Let  $g \in \mathbb{K}[x]$  is a non-zero polynomial. Every polynomial  $f \in \mathbb{K}[x]$  can be written uniquely as:*

$$f = qg + r$$

where  $q, r \in \mathbb{K}[x]$ , and either  $r = 0$  or  $\deg(r) < \deg(g)$ .

While this theorem might seem innocuous when first learned as a teenager, it turns out to be very powerful when studying ideals and polynomials in  $\mathbb{K}[x]$ . The proof of this theorem also provides an algorithm for computing  $q$  and  $r$ , which presented in pseudocode looks something like

---

**Algorithm 1** Division Algorithm

---

**Require:**  $g, f$

**Ensure:**  $q, r$

$q \leftarrow 0; r \leftarrow f$

**while**  $r \neq 0$  **and**  $\text{LT}(g)$  divides  $\text{LT}(r)$  **do**

$q \leftarrow q + \text{LT}(r)/\text{LT}(g)$

$r \leftarrow r - (\text{LT}(r)/\text{LT}(g))g$

**end while**

---

The goal of this worksheet is begin exploring how we might extend this theorem to polynomial rings in more variables. Along the way we will see how these explorations will lead to both computational tools for studying the commutative algebra of polynomial rings over fields, but also give us useful tools for proving theorems in this setting as well.

- (1) Using Theorem 1 to prove the following:
  - (a) If  $f \in \mathbb{K}[x]$  is a non-zero polynomial then  $f$  has at most  $\deg(f)$  roots in  $\mathbb{K}$ .
  - (b) If  $g \in \mathbb{K}[x]$  is a non-zero polynomial then  $f \in \langle g \rangle$  if and only if  $r = 0$ .
  
- (2) Implement algorithm above in Macaulay2, use your code to compute  $q$  and  $r$  for the following polynomials. (If you want to test your code verse the built-in functions use `time f//g` and `time f%g` to compute  $q$  and  $r$  respectively and time the computation.)
  - (a)  $f = x^3 + x + 1, \quad g = x + 1.$
  - (b)  $f = x^5 - 1, \quad g = x - 1.$
  - (c)  $f = x^2 + 1, \quad g = x^3 + x.$
  - (d)  $f = x^{100} - 1, \quad g = x^2 - 1.$
  - (e)  $f = x^{1000} - 1, \quad g = x - 1.$
  - (f)  $f = \sum_{i=0}^{500} x^i, \quad g = x^2 + x + 1.$

**Monomial Orderings.** A monomial ordering on  $\mathbb{K}[x_1, \dots, x_n]$  is a relation  $<$  on  $\mathbb{Z}_{\geq 0}^n$  satisfying the following conditions:

- (i)  $<$  is a total ordering on  $\mathbb{Z}_{\geq 0}^n$ ,
- (ii) if  $\alpha < \beta$  and  $\gamma \in \mathbb{Z}_{\geq 0}^n$  then  $\alpha + \gamma < \beta + \gamma$ , and
- (iii)  $<$  is a well-ordering on  $\mathbb{Z}_{\geq 0}^n$ .

Recall that a *total ordering* means that of any two elements  $\alpha, \beta \in \mathbb{Z}_{\geq 0}^n$  exactly one of the following statements is true:  $\alpha < \beta$ ,  $\alpha = \beta$ , or  $\beta < \alpha$ . That  $<$  is a well-ordering on  $\mathbb{Z}_{\geq 0}^n$  means that every non-empty subset of  $\mathbb{Z}_{\geq 0}^n$  has a smallest element with respect to  $<$ .

- (3) Prove that a monomial ordering  $<$  on  $\mathbb{Z}_{\geq 0}^n$  is equivalent to a relation  $<$  on the set of monomials in  $\mathbb{K}[x_1, \dots, x_n]$  satisfying analogous conditions to (i), (ii), and (iii) above. Explain why this justifies the moniker “monomial ordering”.
- (4) Prove or disprove that the following are monomial orderings:
  - (a) Lexicographic (Lex) Order: Given  $\alpha = (\alpha_1, \dots, \alpha_n)$  and  $\beta = (\beta_1, \dots, \beta_n)$  in  $\mathbb{Z}_{\geq 0}^n$  we say  $\beta <_{\text{lex}} \alpha$  if and only if the left-most non-zero entry of  $\alpha - \beta$  is positive.
  - (b) Graded Lex Order: With  $\alpha$  and  $\beta$  as in (a) we say that  $\beta <_{\text{grlex}} \alpha$  if and only if

$$|\beta| := \sum_{i=1}^n \beta_i < |\alpha| = \sum_{i=1}^n \alpha_i \quad \text{or} \quad |\beta| = |\alpha| \text{ and } \beta <_{\text{lex}} \alpha.$$

- (c) Reverse Lex (RevLex) Order: With  $\alpha$  and  $\beta$  as in (a) we say that  $\beta <_{\text{revlex}} \alpha$  if and only if the right-most non-zero entry of  $\alpha - \beta$  is negative.
- (d) Graded RevLex Order: With  $\alpha$  and  $\beta$  as in (a) we say that  $\beta <_{\text{grevlex}} \alpha$  if and only if

$$|\beta| := \sum_{i=1}^n \beta_i < |\alpha| = \sum_{i=1}^n \alpha_i \quad \text{or} \quad |\beta| = |\alpha| \text{ and } \beta <_{\text{revlex}} \alpha.$$

- (e) Weight Order: Fix a vector  $w = (w_1, \dots, w_n) \in \mathbb{Z}^n$ . With  $\alpha$  and  $\beta$  as in (a) we say that  $\beta <_w \alpha$  if and only if

$$\beta \cdot w = \sum_{i=1}^n \beta_i w_i < \alpha \cdot w = \sum_{i=1}^n \alpha_i w_i$$

- (5) **Leading Terms.** Fix a monomial ordering  $<$  on  $\mathbb{Z}_{\geq 0}^n$ , and use the corresponding ordering on monomials in  $\mathbb{K}[x_1, \dots, x_n]$ . Let

$$f = \sum_{\alpha} c_{\alpha} x^{\alpha} \in \mathbb{K}[x_1, \dots, x_n]$$

be a non-zero polynomial. Define the *leading monomial*  $\text{LM}(f)$ , the *leading coefficient*  $\text{LC}(f)$ , and the *leading term*  $\text{LT}(f)$  with respect to the fixed monomial ordering.

- (a) Compute  $\text{LM}(f)$ ,  $\text{LC}(f)$ , and  $\text{LT}(f)$  for

$$f = x^3 + 3x^2y + 5xy^3 - 7z^2 \in \mathbb{K}[x, y, z]$$

with respect to Lex, Graded Lex, and Graded RevLex orderings (assume  $x > y > z$ ).

- (b) Prove that if  $f, g \in \mathbb{K}[x_1, \dots, x_n]$  are non-zero, then

$$\text{LM}(fg) = \text{LM}(f)\text{LM}(g) \quad \text{and} \quad \text{LT}(fg) = \text{LT}(f)\text{LT}(g).$$

- (c) Explain which property of a monomial ordering is used in part (b).

- (6) Let  $G = (g_1, \dots, g_s)$  be an ordered list of non-zero polynomials in  $\mathbb{K}[x_1, \dots, x_n]$ .
- (a) Write pseudocode for a division algorithm which takes as input  $f$  and  $G$  and returns polynomials  $q_1, \dots, q_s, r$  such that

$$f = q_1g_1 + \dots + q_sg_s + r,$$

where no monomial appearing in  $r$  is divisible by any of the monomials  $\text{LM}(g_1), \dots, \text{LM}(g_s)$ .

- (b) In the algorithm above, what should happen if the leading monomial of the current polynomial is not divisible by any  $\text{LM}(g_i)$ ?
- (c) Prove that your algorithm terminates.
- (d) What plays the role of “degree” in the proof of termination?
- (7) Work in  $\mathbb{K}[x, y]$  with Lex order and  $x > y$ . Let

$$f = xy^2 - x, \quad g_1 = xy - 1, \quad g_2 = y^2 - 1.$$

- (a) Divide  $f$  by the ordered list  $(g_1, g_2)$ .
- (b) Divide  $f$  by the ordered list  $(g_2, g_1)$ .
- (c) Show directly that  $f \in \langle g_1, g_2 \rangle$ .
- (d) What do your computations show about division in several variables? In particular, is the remainder uniquely determined by the ideal  $\langle g_1, g_2 \rangle$ ?